

5

WHAT IS CLAIMED IS:

1. A pharmaceutical kit for nasal drug delivery comprising:
an aqueous solution of cyanocobalamin and excipients in a container and;
a droplet-generating actuator attached to said container and fluidly connected to the
10 cyanocobalamin solution in the container;
wherein said actuator produces a spray of the cyanocobalamin solution through a tip of
the actuator when said actuator is engaged, wherein said spray of cyanocobalamin
solution has a spray pattern ellipticity ratio of from about 1.0 to about 1.4 when measured
at a height of 3.0 cm from the actuator tip.
- 15
2. The kit of claim 1 wherein said spray comprises droplets wherein less than 5% of said
droplets are less than 10 μm in size.
- 20
3. The kit of claim 1 wherein in the aqueous solution of cyanocobalamin has a viscosity
of less than 1000 cPs, and wherein said solution of cyanocobalamin has a bioavailability
of cyanocobalamin of about 7% relative to an intramuscular injection of cyanocobalamin,
with the proviso that mercury and mercury containing compounds are not present in the
solution.
- 25
4. The kit of claim 3 wherein the spray is comprised of droplets of the cyanocobalamin
solution wherein less than 5% of the droplets are less than 10 μm in size.
- 30
5. The kit of claim 3 wherein the spray has a spray pattern major axis and minor axis of
between 25 and 40 mm each.
6. The kit of claim 3 wherein the solution of cyanocobalamin is further comprised of
citric acid, and sodium citrate wherein the solution has a pH of from about 4-6.

- 5 7. The kit of claim 6 wherein the pH of the solution is about 5.
8. The kit of claim 3 wherein cyanocobalamin is present in solution at a concentration of between 0.5-1% by weight.
- 10 9. The kit of claim 8 wherein the concentration of cyanocobalamin in solution is about 0.5%.
10. The kit of claim 6 wherein the citric acid is present in solution at a concentration of about 0.12%, and the sodium citrate is present in solution at a concentration of about
15 0.32%, in water.
11. The kit of claim 3 wherein the cyanocobalamin spray is comprised of droplets of the cyanocobalamin solution wherein 50% of the droplets are 26.9 μm or less in size.
- 20 12. The kit of claim 3 wherein the cyanocobalamin spray is comprised of droplets of the cyanocobalamin solution, wherein 90% of the droplets are 55.3 μm or less in size.
13. The product of claim 3 wherein the cyanocobalamin spray produces droplets of the solution, and wherein 10% of the droplets are 12.5 μm or less in size.
25
14. A kit for administering intranasally a cyanocobalamin solution comprised of a container, a solution of cyanocobalamin in the container, and an actuator attached to said container, wherein a spray of cyanocobalamin solution is expelled through a tip of said actuator when said actuator is engaged wherein said aqueous solution of cyanocobalamin
30 is comprised of cyanocobalamin at a concentration of about 0.5% of total weight of solution, citric acid at a concentration of about 0.12%, sodium citrate at a concentration of about 0.32%, glycerin at a concentration of about 2.23%, benzalkonium chloride at concentration of about 0.02% and water wherein said solution of cyanocobalamin is suitable for intranasal administration, has a viscosity less than about 1000 cPs, and

5 wherein said solution of cyanocobalamin has a bioavailability of cyanocobalamin when
administered intranasally of at least about 7% relative to an intramuscular injection of
cyanocobalamin, with the proviso that the solution of cyanocobalamin contains no
mercury or mercury-containing compounds, and wherein the spray has a spray pattern
ellipticity ratio of from about 1.0 to about 1.4 when measured at a height of 3.0 cm from
10 the actuator tip.

15 15. The kit of claim 14 wherein the cyanocobalamin spray is comprised of droplets of the
cyanocobalamin solution, wherein less than 5 % of the droplets of the cyanocobalamin
spray are less than 10 μm in size.

16. The kit of claim 12 wherein the cyanocobalamin spray is comprised of droplets of the
cyanocobalamin solution, and wherein 50% of the droplets of the cyanocobalamin spray
are 26.9 μm or less in size.

20 15. The method of claim 12 wherein the cyanocobalamin spray produces droplets of the
solution, and wherein 90% of the droplets are 55.3 μm or less in size.

16. The method of claim 12 wherein the cyanocobalamin spray produces droplets of the
solution, and wherein 10% of the droplets are 12.5 μm or less in size.

25 17. The method of claim 12 wherein the spray has a spray pattern major axis of about
35.3 mm and a minor axis of about 30.8 mm.

30 18. A method for administering cyanocobalamin intranasally comprised of providing an
aqueous solution of cyanocobalamin, wherein the solution of cyanocobalamin has a
viscosity of less than 1000 cPs, and wherein said solution of cyanocobalamin has a
bioavailability of cyanocobalamin of about 7% relative to an intramuscular injection of
cyanocobalamin, with the proviso that mercury and mercury containing compounds are
not present in the solution, wherein the cyanocobalamin formulation is administered into

- 5 a nose of an individual through an actuator tip as a spray, wherein the spray has a spray pattern ellipticity ratio of from about 1.0 to about 1.4 when measured at a height of 3.0 cm from the actuator tip.
19. The method of claim 18 wherein the spray produces droplets, wherein less than 5%
10 of the droplets are less than 10 μm in size.
20. The method of claim 18 wherein the spray has a spray pattern major axis and minor axis of between 25 and 40 mm each.
- 15 21. The method of claim 18 wherein the solution of cyanocobalamin is further comprised of citric acid, and sodium citrate wherein the solution has a pH of from about 4-6.
22. The method of claim 21 wherein the pH of the solution is about 5.
- 20 23. The method of claim 18 wherein cyanocobalamin is present in solution at a concentration of between 0.5-1% by weight.
24. The method of claim 6 wherein the concentration of cyanocobalamin in solution is about 0.5%.
- 25 25. The method of claim 18 wherein the citric acid is present in solution at a concentration of about 0.12%, and the sodium citrate is present in solution at a concentration of about 0.32%, in water.
- 30 26. The method of claim 18 wherein the cyanocobalamin spray produces droplets of the solution, and wherein 50% of the droplets are 26.9 μm or less in size.

5 27. The method of claim 18 wherein the cyanocobalamin spray produces droplets of the solution, and wherein 90% of the droplets are 55.3 μm or less in size.

28. The method of claim 18 wherein the cyanocobalamin spray produces droplets of the solution, and wherein 10% of the droplets are 12.5 μm or less in size.

10

29. A method for administering cyanocobalamin comprised of providing an aqueous solution of cyanocobalamin wherein said aqueous solution of cyanocobalamin is comprised of cyanocobalamin at a concentration of about 0.5% of total weight of solution, citric acid at a concentration of about 0.12%, sodium citrate at a concentration of about 0.32%, glycerin at a concentration of about 2.23%, benzalkonium chloride at concentration of about 0.02% and water wherein said solution of cyanocobalamin is suitable for intranasal administration, has a viscosity less than about 1000 cPs, and wherein said solution of cyanocobalamin has a bioavailability of cyanocobalamin when administered intranasally of at least about 7% relative to an intramuscular injection of cyanocobalamin, with the proviso that the solution of cyanocobalamin contains no mercury or mercury-containing compounds, and wherein the cyanocobalamin solution is administered into a nose of an individual through an actuator tip as a spray, wherein the spray has a spray pattern ellipticity ratio of from about 1.0 to about 1.4 when measured at a height of 3.0 cm from the actuator tip.

25

30. The method of claim 29 wherein the cyanocobalamin spray produces droplets of the solution, wherein less than 5 % of the droplets are less than 10 μm in size.

31. The method of claim 29 wherein the cyanocobalamin spray produces droplets of the solution, and wherein 50% of the droplets are 26.9 μm or less in size.

30

32. The method of claim 29 wherein the cyanocobalamin spray produces droplets of the solution, and wherein 90% of the droplets are 55.3 μm or less in size.

5 33. The method of claim 29 wherein the cyanocobalamin spray produces droplets of the solution, and wherein 10% of the droplets are 12.5 μm or less in size.

34. The method of claim 29 wherein the spray has a spray pattern major axis and a minor axis of about 25 - 40 mm each.

10

35. A method for elevating the vitamin B12 levels in the cerebral spinal fluid (CSF) comprising administering intranasally a sufficient amount of a solution of cyanocobalamin so that the average ratio of vitamin B12 in the CSF to that in the blood serum ($\text{B12 CSF/B12 Serum} \times 100$) is increased to at least about 1.1, wherein said
15 aqueous solution of cyanocobalamin is comprised of cyanocobalamin at a concentration of about 0.5% of total weight of solution, citric acid at a concentration of about 0.12%, sodium citrate at a concentration of about 0.32%, glycerin at a concentration of about 2.23%, benzalkonium chloride at concentration of about 0.02% and water wherein said solution of cyanocobalamin is suitable for intranasal administration, has a viscosity less
20 than about 1000 cPs, and wherein said solution of cyanocobalamin has a bioavailability of cyanocobalamin when administered intranasally of at least about 7% relative to an intramuscular injection of cyanocobalamin, with the proviso that the cyanocobalamin solution contains no mercury or mercury-containing compounds and wherein the cyanocobalamin solution is administered into a nose of an individual through an actuator
25 tip as a spray, wherein the spray has a spray pattern ellipticity ratio of from about 1.0 to about 1.4 when measured at a height of 3.0 cm from the actuator tip.

36. The method of claim 35 wherein the cyanocobalamin spray produces droplets of the solution, wherein less than 5 % of the droplets are less than 10 μm in size.

30

37. The method of claim 35 wherein the cyanocobalamin spray produces droplets of the solution, and wherein 50% of the droplets are 26.9 μm or less in size.

38. The method of claim 35 wherein the cyanocobalamin spray produces droplets of the
35 solution, and wherein 90% of the droplets are 55.3 μm or less in size.

5

39. The method of claim 35 wherein the cyanocobalamin spray produces droplets of the solution, and wherein 10% of the droplets are 12.5 μm or less in size.

40. The method of claim 35 wherein the spray has a spray pattern major axis and a minor
10 axis of between 25 - 40 mm each.